

The 2013 Multi-Modality Appropriate Use Criteria for the Detection and Risk Assessment of Stable Ischemic Heart Disease document replaces the following patient indications from previously published appropriate use criteria documents:

- Symptomatic - Table 1.1, Indications 1-6
 - RNI (2009) – Table 1, Indications 1-5
 - Echo (2011) – Table 9, Indications 114-118
 - CTA (2010) – Table 1, Indications 1-8
 - MR (2006) – Table 12, Indications 1-7
 - Diagnostic Cath (2012) – Table 1.2, Indications 8-10
 -
- Asymptomatic
 - RNI (2009) – Table 2, Indications 12-15
 - Echo (2011) – Table 10, Indications 124-127
 - CTA (2010) – Table 1, Indication 7
 - Diagnostic Cath (2012) – Table 1.2, Indications 5-7
- Prior Testing
 - RNI (2009) – Table 3, Indications 23 - 26, 33-36
 - Echo (2011) – Table 12, Indications 136-139,142-145
 - CTA (2010) – Table 4, Indications 27-28
 - MR (2006) – Table 13, Indications 11-13
 - Diagnostic Cath (2012) – Indications 11–17, 19–21, 28-39
- Prior Procedure
 - RNI (2009) – Table 6 Indications 55-60
 - Echo (2011) – Table 15 Indications 169-174
 - CTA (2010) – Table 6, Indication 40, 42, 44
 - MR (2006) – Table 15, Indications 16-17
 - Diagnostic Cath (2012) – Table 1.5, Indications 52-56
- Preoperative
 - RNI (2009) – Table 4, Indications 40-48
 - Echo (2011) – Table 13, Indications 154-162
 - CTA (2010) – Table 5, Indications 30-38
 - MR (2006) – Table 14, Indications 14-15
 - Diagnostic Cath (2012) – Table 1.7, Indications 64-69
- Rehab Evaluation
 - RNI (2009) – Table 6, Indication 61
 - Echo (2011) – Table 14, Indication 168